arXiv:1008.3674v1 [math.NT] 22 Aug 2010

EXPLICIT FACTORIZATION OF PRIME INTEGERS IN QUARTIC NUMBER FIELDS DEFINED BY X^4+aX+b

L HOUSSAIN EL FADIL

Abstract. For every prime integer p, an explicit factorization of the principal ideal $p\mathbb{Z}_K$ into prime ideals of $\mathbb{Z}_K$ is given, where K is a quartic number field defined by an irreducible polynomial $X^4+aX+b \in \mathbb{Z}[X]$.

Key words: Prime ideal factorization, Newton polygons, Quartic number fields.
AMS classification: 11Y40.

Introduction

Let K be a quartic number field defined by an irreducible polynomial $P(X)=X^4+aX+b \in \mathbb{Z}[X]$, $\mathbb{Z}_K$ its ring of integers, $\triangle$ the discriminant of P, d_K its discriminant, α a complex root of $P(X)$ and $ind(P)=[\mathbb{Z}_K:\mathbb{Z}[\alpha]]$ the index of $\mathbb{Z}[\alpha]$. In this paper, the goal is to give an explicit factorization of $p\mathbb{Z}_K$ into prime ideals of $\mathbb{Z}_K$; the form $p\mathbb{Z}_K=\prod_{i=1}^r P_i^{e_i}$ and for every prime factor P_i, an integral element w_i such that $P_i=(p,w_i)$ are given. Let p be a prime integer. It is well known that if p does not divide $ind(P)$, then the Dedekind's theorem gives us an explicit factorization of the principal ideal $p\mathbb{Z}_K$ into prime ideals of $\mathbb{Z}_K$: $p\mathbb{Z}_K$ is p-analogous to the factorization of $\bar{P}(X)$ modulo p. (see, for example [3, page 257]).

Theorem 0.1. *Let p be a prime integer. Denote by $^-$ the canonical map of $\mathbb{Z}[X]$ into $F_p[X]$, and let $\bar{P}(X)=\prod_{i=1}^r g_i(X)^{e_i}$, where $g_1(X)$,..,$g_r(X)$ are distinct irreducible in $F_p[X]$ and e_1,..e_r are positive integers. For every i, let $P_i=(p,f_i(\alpha))$, where $f_i \in \mathbb{Z}[X]$ is a monic lifting over $g_i(X)$. Then*
If p does not divides $ind(P)$, then $p\mathbb{Z}_K=\prod_{i=1}^r P_i^{e_i}$ and for every i, $e(P_i/p)=e_i$ and $f(P_i/p)=f_i=deg(g_i)$.

If p is not a common index divisor of K, then there exist an element $\phi \in \mathbb{Z}_K$ which generates $\mathbb{Z}_K$ and $v_p(ind(\pi_\phi)=0$, where π_ϕ is the minimal polynomial of ϕ, and then we can apply Dedekind's theorem to obtain the prime ideal decomposition explicitly. However given α, it is not easy to determine such an element ϕ in general. The construction of ϕ was based on the p-integral bases of $\mathbb{Z}_K$ given in [1]. If p is common index divisor of K, then for every prime P factor of $p\mathbb{Z}_K$, an element $\beta \in \mathbb{Z}_K$ such that $v_P(\beta)=1$ and for every prime ideal $Q \neq P$ above p, $v_Q(\beta)=0$ will be constructed. A such $\beta \in \mathbb{Z}_K$ satisfies $P=(p,\beta)$.

For every prime p, let v_p be the p-adic discrete valuation defined in $\mathbb{Q}_p$ by $v_p(p)=1$. v_p is extended to $\mathbb{Q}_p[X]$ by $v_p(A(X)) := Min\{v_p(a_i)\,,\, 0 \le i \le r\}$, where $A(X)=\sum_{i=0}^r a_i X^i$. For every $(x,m) \in \mathbb{Z}^2$, denote $x_p=\frac{x}{p^{v_p(x)}}$, $x \pmod m$ the remainder of the Euclidean division of x by m. For every odd prime p, denote $(\frac{-}{p})$ the Legendre symbol. If p does not divide x and there exists $t \in \mathbb{Z}$ such that $t^n=x \pmod p$, then let $(\frac{x}{p})_n=1$. Otherwise, $(\frac{x}{p})_n \neq 1$.

1. Newton polygons

Let $F(X)$ be an irreducible polynomial in $\mathbb{Z}[X]$, $\phi \in \mathbb{Z}[X]$ a monic polynomial of degree at least 1. Let $F(X)=\sum_{i=0}^n a_i(X)\phi(X)^i = a_0(X)+..+a_{n-1}(X)\phi(X)^{n-1}+a_n(X)\phi(X)^n$ be

---

Supported by ERCIM.

the ϕ-adic development of $F(X)$ (for every i, $deg(a_i(X)) \le deg(\phi) - 1$). Let p be a prime integer. The ϕ-Newton polygon of $F(x)$, with respect to p, is the lower convex envelope of the set of points (i, u_i), $u_i < \infty$, in the Euclidian plane, where $u_i = v_p(a_i(X))$.



The ϕ-Newton polygon is the union of different adjacent sides $S_0, \dots, S_g$ with increasing slope $\lambda_0 < \lambda_1 < \cdots < \lambda_g$. We shall write $N_\phi(F) = S_0 + \cdots + S_g$. The principal part of $N_\phi(F)$, denoted $N_\phi^+(F)$, is the polygon determined by the sides of negative slope of $N_\phi(F)$.

For every $0 \le i \le n$, we attach to any abscissa the following residual coefficient $c_i \in \mathbb{F}_\phi$:

$$c_i = \begin{cases} 0, & \text{if } (i, u_i) \text{ lies strictly above } N \text{ or } u_i = \infty, \\ \text{red}\left(\dfrac{a_i(X)}{p^{u_i}}\right), & \text{if } (i, u_i) \text{ lies on } N. \end{cases}$$

Let S be one of the sides of N, with slope λ, and let $\lambda = -h/e$, with h, e positive coprime integers. Let $l = \ell(S)$ be the length of the projection of S to the x-axis, $d(S) := \ell(S)/e$: the degree of S. Note that S is divided into $d(S)$ segments by the points of integer coordinates that lie on S. Let s be the initial abscissa of S and d the degree of S. If ϕ is a monic polynomial such that $\bar{\phi}$ is irreducible factor of $\bar{F}(X)$ modulo p, then let $\mathbb{F}_\phi := \frac{\mathbb{F}_p[X]}{(\bar{\phi})}$ and $F_S(Y) := c_s + c_{s+e}Y + \cdots + c_{s+(d-1)e}Y^{d-1} + c_{s+de}Y^d \in \mathbb{F}_\phi[Y]$, the residual polynomial of $f(X)$ attached to S.

In the remainder, $F(X)$ is a monic irreducible polynomial, θ is a complex root of $F(X)$ and ϕ is a monic polynomial such that $\bar{\phi}$ is irreducible factor of $\bar{F}(X)$ modulo p. Let $N = S_k + .. + S_1 + S_0$ be the $X\phi$-Newton polygon of $F(X)$ with respective slopes $\lambda_k < .. < \lambda_1 < \lambda_0$. $F(X)$ is said to be ϕ-regular if for every i, $F_{S_i}(Y)$ is square free.

2. Main results

In this section, for every prime integer p, an explicit factorization of the principal ideal $p\mathbb{Z}_K$ into prime ideals of $\mathbb{Z}_K$ is given (Note only the form is given, but two element generators of each prime ideal factor are given). Recall the celebrated Theorem of Hensel:

Theorem 2.1 (Hensel)**.** *Let $P(X) \in \mathbb{Z}[X]$ be a monic irreducible polynomial, p a prime integer, $\mathbb{Q}_p$ the p-adic completion of $\mathbb{Q}$ and $\mathbb{Z}_p$ its ring of integers. Let $P(X) = F_1(X) \cdots F_t(X)$ be the factorization into a product of monic irreducible polynomials in $\mathbb{Q}_p[X]$ and consider the local fields $K_i = \mathbb{Q}_p[X]/(F_i(X))$, for $i = 1, \dots, t$. The factorization of $p\mathbb{Z}_K$ into a product of prime ideals of $\mathbb{Z}_K$ is: $p\mathbb{Z}_K = \mathfrak{p}_1^{e_1} \cdots \mathfrak{p}_t^{e_t}$, where $f(\mathfrak{p}_i) = f(K_i/\mathbb{Q}_p)$ and $e_i = e(K_i/\mathbb{Q}_p)$.*

Lemma 2.2. *Assume that $P(X) = \bar{\phi}^l(X) \pmod{p}$ such that $N_\phi P = S$ is one side and $P_S(Y)$ is irreducible in $\mathbb{F}_\phi[Y]$, where $\mathbb{F}_\phi = \frac{\mathbb{F}_p[X]}{(\bar{\phi}(X))}$. Let $\lambda = -h/e$ be the slope of S such that h and e are positive coprime integers. Define $\alpha_S := \phi(\alpha)^e/p^h$. Then $v_p(\alpha_S) = 0$ and $P_S(\alpha_S) = 0$.*

Proof. Let $P(X)=\sum_{k=0}^{l}a_k(X)\phi(X)^k$ be the ϕ-adic development of $P(X)$. Since $N_\phi(P)$ is one side of slope $-\lambda$, we have for every i, $v(a_i(X))\geq(i-l)\lambda$ and $v(a_0(X))=-l\lambda$. Thus, $v_p(\phi(\alpha))=-\lambda$. Moreover, since for every $0\leq j\leq d$ and for every $je<k<(j+1)e$, red $\frac{a_k(X)}{p^{u_k}}=0$ and $\frac{P(\alpha)}{p^l}=0$, then $\overline{(\frac{\phi(\alpha)^e}{p^h})^d+\frac{a_{l-e}(\alpha)}{p^h}\frac{\phi(\alpha)^e}{p^h})^{d-1}+\cdots+\frac{a_e(\alpha)}{p^{h(d-1)}}\frac{\phi(\alpha)^e}{p^h})+\frac{a_{le}(\alpha)}{p^l}}=0$ in $\mathbb{F}_\phi$. Therefore, $P_S(\alpha_S))=0$ in $\mathbb{F}_\phi$. □

Let $P(X)\in\mathbb{Z}[X]$ be a monic irreducible polynomial and $\bar{P}(X)=\prod_{i=1}^{r}\phi_i^{l_i}\pmod{p}$ its factorization into irreducible polynomials of $\mathbb{F}_p[X]$. For every $i:=1..r$, let $N_i:=N_{\phi_i}^{+}(P)=S_1^i+\cdots+S_{k_i}^i$ and for every $j:=1..k_i$, let $P_{S_j}(Y)=\prod_{s=1}^{r_{ij}}\psi_s(Y)^{n_s}$ be the factorization of $P_{S_j}(Y)$ into irreducible polynomials of $\mathbb{F}_\phi[Y]$, where $\mathbb{F}_\phi=\frac{\mathbb{F}_p[X]}{(\phi(X))}$.

Theorem 2.3. *Under these hypothesis, if every $P_{S_j}(Y)$ is square free (every $n_s=1$), then*

$$p\mathbb{Z}_K=\prod_{i=1}^{r}\prod_{j=1}^{k_i}\prod_{s=1}^{r_{ij}}\mathfrak{p}_s^{e(S_j^i)},$$

and for every $i:=1..r$, $j:=1..k_i$ and $s:=1..r_{ij}$, $v_{\mathfrak{p}_s}(\phi_i(\alpha))=-e(S_j^i)\lambda_j$, where $e(S_j^i)$ is the ramification index of the side S_j^i and $-\lambda_j$ is its slope.

Proof. By Hensel Theorem, it suffices to factorize $P(X)$ into irreducible polynomials of $\mathbb{Q}[X]$. By Hensel Lemma, it suffices to show this result for $r=1$. By Theorem of the polygon we can assume that $N_1=S$ is one side. Since $P_S(Y)$ is square free, by Theorem of the residual polynomial, we can assume that $P_S(Y)$ is irreducible in $\mathbb{F}_\phi[Y]$. By Theorem of the product, if $P_S(Y)$ is irreducible in $\mathbb{F}_\phi[Y]$, then $P(X)$ is irreducible in $\mathbb{Q}[X]$.

Now, assume that $P(X)\in\mathbb{Z}_p[X]$ is a monic irreducible polynomial in $\mathbb{Q}[X]$ such that $\bar{P}(X)=\phi^l(X)\pmod{p}$, $N_\phi(P)=S$ is one side and $P_S(Y)$ is irreducible in $\mathbb{F}_\phi[Y]$ of degree $d(S)$. Denote $\mathbb{K}=\mathbb{Q}_p[\alpha]$, $\mathbb{Z}_{\mathbb{K}}$ its ring of integers over $\mathbb{Z}_p$ and $\mathfrak{p}$ the maximal ideal of $\mathbb{Z}_{\mathbb{K}}$. By Hensel Theorem, $p\mathbb{Z}_{\mathbb{K}}=\mathfrak{p}^{e(\mathfrak{p})}$, where $e(\mathfrak{p})=\frac{deg(P)}{f(\mathfrak{p})}$.

Let $\lambda=-\frac{e}{h}$ be the slope of S such e and h are positive coprime. Let $\alpha_S=\frac{\phi^e(\alpha)}{p^h}$ and $\iota\colon\mathbb{Z}_p[X,Y]\longrightarrow\mathbb{Z}_p,\quad a(X,Y)\mapsto a(\alpha,\alpha_S)$. Then the ideal $\mathfrak{m}=\iota^{-1}(\mathfrak{p})$ is a maximal ideal of $\mathbb{Z}_p[X,Y]$ generated by $(p,\phi(X),\varphi(X,Y))$, where $\varphi(X,Y)\in\mathbb{Z}_p[X,Y]$ is a monic polynomial such that $\psi(Y):=\mathrm{red}(\varphi(X,Y))$ is irreducible in $\mathbb{F}_\phi[Y]$. Thus, $\frac{\mathbb{Z}_{\mathbb{K}}}{\mathfrak{p}}\simeq\frac{\mathbb{F}_\phi[Y]}{(\psi(Y))}$ and $f(\mathfrak{p})=d(S).m$. Thus, $e(\mathfrak{p})=\frac{deg(P)}{d(S).m}=\frac{l.m}{d(S).m}=e(S)$, where $m=deg(\phi)$.
On the other hand, since $N_1=S$ is one side of slope λ, we have $v_p(\phi(\alpha)^l)=v_p(a_0(\alpha))=-l\lambda$, $v_p(\phi(\alpha))=-\lambda$ and $v_{\mathfrak{p}}(\phi(\alpha))=-e(\mathfrak{p})\lambda$. □

Let $P(X)=X^4+aX+b$. Note that if there exists a prime p such that $v_p(a)\geq 3$ and $v_p(b)\geq 4$, then let q_1 and q_2 be respectively the quotient of $v_p(a)$ by 3 and of $v_p(b)$ by 4. Let $\theta:=\frac{\alpha}{p^q}$, where $q:=Min(q_1,q_2)$. Then θ is integral with minimal polynomial $F(X)=X^4+AX+B\in\mathbb{Z}[X]$, where $A=\frac{a}{p^{3q}}$ and $B=\frac{b}{p^{4q}}$. As $K=\mathbb{Q}[\theta]=\mathbb{Q}[\alpha]$, then up to replace α by $\frac{\alpha}{p^q}$, we can assume that for every prime p, $v_p(a)\leq 2$ or $v_p(b)\leq 3$.

Theorem 2.4. *Let p be a prime integer. In the following tables, the form of $p\mathbb{Z}_K$ as a product of prime ideals of $\mathbb{Z}_K$, and for every prime factor P of $p\mathbb{Z}_K$ an integral element ϕ of K such that $P=(p,\phi)$ are given: If $p\mathbb{Z}_K=\prod_i P_i^{e_i}$, then for every $i\neq j$, an element $\beta_i\in K$ such that $v_{P_i}=(\beta_i)=1$ and $v_{P_j}=(\beta_i)=0$ is given.*

$$TableA: v_p(a)\geq 1\, and\, v_p(b)\geq 1$$

Case	*Conditions*	p	$p\mathbb{Z}_K$	*Generators*
A1	$v_p(b) = 3,\ v_p(a) \geq 3$		P^4	$P = (p, \frac{\alpha^3}{p^2})$
A2	$v_p(b) \geq 3,\ v_p(a) = 2$		$P_1P_2^3$	$P_1 = (p, \frac{\alpha^3}{p^2}),\ P_2 = (p, a_p + \frac{\alpha^3}{p^2})$
A3	$v_p(b) \geq 2,\ v_p(a) = 1$		$P_1P_2^3$	$P_1 = (p, \frac{\alpha^3}{p}),\ P_2 = (p, a_p + \frac{\alpha^3}{p})$
A4	$v_p(b) = 2,\ v_p(a) \geq 2$ $(\frac{-b_p}{p}) = -1$	$\neq 2$	P^2	$P = (p, \frac{\alpha^3}{p})$
A5	$v_p(b) = 2,\ v_p(a) = 2$ $(\frac{-b_p}{p}) = 1$	$\neq 2$	$P_1^2P_2^2$	$P_1 = (p, t + \frac{\alpha^2}{p})$ $P_2 = (p, -t + \frac{\alpha^2}{p}),\ (v_p(t^2 + b_p) = 1)$
A6	$v_p(b) = 2,\ v_p(a) \geq 3$ $(\frac{-b_p}{p}) = 1$	$\neq 2$	$P_1^2P_2^2$	$P_2 = (p, t + \frac{\alpha^3+\alpha^2}{p})$ $P_2 = (p, -t + \frac{\alpha^3+\alpha^2}{p}),\ (t^2 + b_p = 0 \pmod{p})$
A7	$v_p(b) = 1,\ v_p(a) \geq 1$		P^4	$P = (p, \alpha)$
A8	$v_p(b) = 2,\ v_p(a) \geq 2$	2		*go to* $TableA8$

$TableA8:\ v_2(b) = 2\, and\, v_2(a) \geq 2$

Case	*conditions*	ϕ_2	$2\mathbb{Z}_K$	*Generators*
A8.1	$v_2(b) = 2,\ v_2(a) = 2$	$\frac{\alpha^2+2}{2}$	P^4	$P = (2, \phi_2)$
A8.2	$b = 12 \pmod{16},\ v_2(a) = 3$	$\frac{\alpha^3+2\alpha}{4}$	P^4	$P = (2, \phi_2)$
A8.3	$b = 4 \pmod{16},\ v_2(a) = 3$	$\frac{\alpha^3+2\alpha^2+2\alpha}{4}$	P^4	$P = (2, \phi_2)$
A8.4	$b = 4 \pmod{32}, v_2(a) = 4$	$\frac{\alpha^2+2\alpha+2}{4}$	P^4	$P = (2, \phi_2)$
A8.5	$b = 20 \pmod{64}, v_2(a) = 4$	$\frac{\alpha^2+2\alpha-2}{4}$	P^2	$P = (2, \phi_2)$
A8.6	$b = 52 \pmod{64}, a = 16 \pmod{64}$	$\frac{\alpha^3-2\alpha^2-2\alpha}{8}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$
A8.7	$b = 52 \pmod{64}, a = 48 \pmod{64}$	$\frac{\alpha^3-2\alpha^2+6\alpha}{8}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$
A8.8	$b = 12 \pmod{32}, v_2(a) \geq 4$	$\frac{\alpha^3+2\alpha}{4}$	P^2	$P = (2, \phi_2)$
A8.9	$b = 28 \pmod{32}, v_2(a) = 4$	$\frac{\alpha^2+2}{4}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$
A8.10	$b = 28 \pmod{32}, v_2(a) \geq 5$	$\frac{\alpha^2+12\alpha+2}{4}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$
A8.11	$b = 20 \pmod{32}, v_2(a) \geq 5$	$\frac{\alpha^2+2\alpha-2}{4}$	P^4	$P = (2, \phi_2)$
A8.12	$b = 36 \pmod{64}, v_2(a) \geq 5$	$\frac{\alpha^2+2\alpha+2}{4}$	P^2	$P = (2, \phi_2)$
A8.13	$b = 4 \pmod{64}, v_2(a) = 5$	$\frac{\alpha^3-2\alpha-4}{8}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$
A8.14	$b = 4 \pmod{64}, v_2(a) \geq 6$	$\frac{\alpha^3+4\alpha^2-2\alpha+4}{8}$	$P_1^2P_2^2$	$P_1 = (2, \phi_2),\ P_2 = (2, \phi_2 + 1)$

$TableB:\ v_p(b) \geq 1\, and\, v_p(a) = 0$

Case	*Conditions*	p	ϕ	$p\mathbb{Z}_K$	*Generators*
B1		2	α	$P_1P_2P_3$ $P_3=(2,1+\alpha+\alpha^2)$	$P_1=(2,\alpha),\ P_2=(2,1+\alpha)$
B2	$(\frac{-a}{p})_3\neq 1$	≥ 5	α	P_1P_2	$P_1=(p,\alpha),\ P_2=(p,a+\alpha^3)$
B3	$(\frac{-3}{p})=-1,\ (\frac{-a}{p})_3=1$	≥ 5	α	$P_1P_2P_3$	$P_1=(p,\alpha),\ P_2=(p,-u+\alpha)$ $P_3=(p,u^2+u\alpha+\alpha^2)$ $u^3=-a \pmod p$
B4	$(\frac{-3}{p})=1,\ (\frac{-a}{p})_3=1$	≥ 5	α	$P_1P_2P_3P_4$	$P_1=(p,\alpha),\ P_2=(p,-u+\alpha)$ $u^3=-a \pmod p,\ P_3=(p,v_1+\alpha)$ $P_4=(p,-(u+v_1)+\alpha)$ $v^2=-3 \pmod p, 2v_1=-u(1+v) \pmod p$
B5	$v_3(b)\geq 2, v_3(a)=0$ $a^2\neq 1 \pmod 9$	*3*	α	$P_1P_2^3$	$P_1=(3,\alpha),\ P_2=(3,\alpha-a)$
B6	$v_3(b)\geq 2,\ a^2=1 \pmod 9$	*3*		$P_1P_2^2P_3$	*go to* $TableB6$
B7	$v_3(a)=0, b=6 \pmod 9$ $a^2\neq 4 \pmod 9$	*3*	α	$P_1^3P_2$	$P_1=(3,\alpha-a),\ P_2=(3,\alpha)$
B8	$b=6 \pmod 9, a^2=4 \pmod 9$	*3*	α	$P_1P_2^2P_3$	*go to* $TableB6$
B9	$v_3(a)=0, b=3 \pmod 9$ $a^2\neq 7 \pmod 9$	*3*	α	$P_1^3P_2$	$P_1=(3,\alpha-a),\ P_2=(3,\alpha)$
B10	$b=3 \pmod 9, a^2=7 \pmod 9$ $v_3(b+a^4-a^2)=2$	*3*		$P_1P_2^3$	$P_1=(3,\theta-4a),\ P_2=(3,\frac{\theta^3-4a\theta^2}{3})$ $\theta=\alpha-a$
B11	$b=3 \pmod 9, a^2=7 \pmod 9$ $v_3(b+a^4-a^2)\geq 3$	*3*			*go to* $TableB11$

$TableB11$

Let $s\in\mathbb{Z}$ *such that* $as=-4b_3 \pmod{3^{v_3(\triangle)+1}}$, *and let* $\theta=\alpha-s,\ A=4s^3+a$ *and* $B=s^4+as+b$.

Case	*Conditions*	ϕ	$3\mathbb{Z}_K$	*Generators*
B11.1	$v_3(\triangle)=6,\ B_3=1 \pmod 3$	$\frac{\theta^3+4s\theta^2+6s^2\theta+A}{9}$	P_1P_2	$P_1=(3,\phi),\ P_2=(3,\phi^3-\phi-s)$
B11.2	$v_3(\triangle)=6$ $B_3=-1 \pmod 3,\ s=1 \pmod 3$	$\frac{\theta^3+4s\theta^2+6s^2\theta+A}{9}$	$P_1P_2P_3$	$P_1=(3,\phi),\ P_2=(3,\phi^2-\phi-1)$ $P_3=(3,\phi+1)$
B11.3	$v_3(\triangle)=6$ $B_3=-1 \pmod 3,\ s=-1 \pmod 3$	$\frac{\theta^3+4s\theta^2+6s^2\theta+A}{9}$	$P_1P_2P_3$	$P_1=(3,\phi),\ P_2=(3,\phi^2+\phi-1)$ $P_3=(3,\phi-1)$
B11.4	*otherwise*		*go to* $TableB11.4$	

$TableC: v_p(b)=0, v_p(a)\geq 1$

	Conditions	p	ϕ	$p\mathbb{Z}_K$	*Generators*
1	$(\frac{2}{p})=1,\ (\frac{b}{p})_4=1$	≥ 5	α	P_1P_2	$P_1=(p,\alpha^2+u\alpha+t),\ t^2=b \pmod p$ $P_2=(p,\alpha^2-u\alpha+t),\ 2t=u^2 \pmod p$
2	$(\frac{-1}{p})=1,(\frac{-b}{p})_4=1$	≥ 5	α	$P_1P_2P_3P_4$	$P_1=(p,\alpha+t),\ P_2=(p,\alpha-t),\ t^4=-b \pmod p$ $P_3=(p,\alpha+ut),\ P_4=(p,\alpha-ut)\ ,\ u^2=-1 \pmod p$
3	$(\frac{-1}{p})=-1,\ (\frac{-b}{p})_4=1$	≥ 5	α	$P_1P_2P_3$	$P_1=(p,\alpha+t),\ P_2=(p,\alpha-t),\ t^4=-b \pmod p$ $P_3=(p,\alpha^2+t^2)$
4	$(\frac{-b}{p})=1,\ (\frac{-b}{p})_4\neq 1$	≥ 5	α	P_1P_2	$P_1=(p,\alpha^2+t),\ P_2=(p,\alpha^2-t),\ t^2=-b \pmod p$
5	$(\frac{4b}{p})_4=1\ ,(\frac{-b}{p})_4\neq 1$	≥ 5	α	P_1P_2	$P_1=(p,\alpha^2+u\alpha+t),\ P_2=(p,\alpha^2-u\alpha+t)$ $u^4=4b \pmod p,\ t^2=b \pmod p$
6	$(\frac{-b}{p})_4\neq 1$ $(\frac{4b}{p})_4\neq 1,\ (\frac{-b}{p})=-1$	≥ 5	α	P	$P=(p)$
7	$b=1 \pmod 3$	*3*	α	P_1P_2	$P_1=(3,\alpha^2+\alpha-1)$ $P_1=(3,\alpha^2-\alpha-1)$
8	$b=-1 \pmod 3$	*3*	α	$P_1^2P_2P_3$	$P_1=(3,\alpha^2+1)$ $P_2=(3,\alpha-1),\ P_3=(3,\alpha+1)$
9	$b=1 \pmod 4,\ a=0 \pmod 4$	*2*	α	P^4	$P=(2,\alpha-1)$
10	$b=3 \pmod 4,\ a=2 \pmod 4$	*2*	α	P^4	$P=(2,\alpha-1)$
11	$b=1 \pmod 4,\ a=2 \pmod 4$	*2*		$P_1^3P_2$	*go to table*8
12	$b=7 \pmod 8,\ a=4 \pmod 8$	*2*		P^2	$P=(2,\alpha-1)$
13	$b=7 \pmod 8,\ a=0 \pmod 8$ $1+b+a=8 \pmod{16}$	*2*	$\frac{\theta^3+4\theta^2+6\theta}{4}$ $\theta=\alpha-1$	$P_1^2P_2$	$P_1=(2,\phi+1)$ $P_2=(2,\phi^2+\phi+1)$
14	$b=7 \pmod 8,\ a=0 \pmod 8$ $1+b+a=0 \pmod{16}$	*2*		$P_1^2P_2P_3$	*go to TableC*14
15	$b=3 \pmod 8,\ a=4 \pmod 8$	*2*		*go to TableC*14	

$$TableD:\ v_p(ab)=0$$

For $p\geq 5$, *let* $s\in\mathbb{Z}$ *such that* $3as+4b=0 \pmod{p^{v_p(\triangle)}}$, *and let* $\theta=\alpha-s$.

Case	*Conditions*	p	ϕ	$p\mathbb{Z}_K$	*Generators*
D1		*2*	α	P	$P=(2)$
D2	$b=-1 \pmod 3$	*3*	α	P	$P=(3)$
D3	$a=b=1 \pmod 3$	*3*	α	P_1P_2	$P_1=(3,\alpha-1),\ P_2=(3,\alpha^3+\alpha^2+\alpha-1)$
D4	$a=b=1 \pmod 3$	*3*	α	P_1P_2	$P_1=(3,\alpha+1),\ P_2=(3,\alpha^3-\alpha^2+\alpha+1)$
D5	$v_p(\triangle)=0$	$p\geq 5$	α	*p-analogous to*$\bar{P}(X)$	
D6	$v_p(\triangle)=1$ $(\frac{-2}{p})=1$	$p\geq 5$	θ	$P_1P_2P_3^2$	$P_3=(p,\theta),\ P_1=(p,\theta-v_1),\ P_2=(p,\theta-v_2)$ $u^2=-2 \pmod p, v_1=-s(2+u), v_2=-s(2-u) \pmod p$
D7	$v_p(\triangle)=1$ $(\frac{-2}{p})=-1$	$p\geq 5$	θ	$P_1P_2^2$	$P_2=(p,\theta),$ $P_1=(p,\theta^2+4s\theta+6s^2)$
D8	$v_p(\triangle)\geq 2$	$p\geq 5$			*go to TableD*8

$$TableB6: B=a^4-a^2+b,\ A=-4a^3+a$$

Case	*Conditions*	$3\mathbb{Z}_K$	β_3	β_2	β_1
B6.1	$v_3(B)=2,\ v_3(A)=1$	$P_1P_2^2P_3$	$\frac{\theta^3-4a\theta^2}{3}$	$\frac{\theta^3-4a\theta^2+6sa^2\theta+A}{3}+\theta^2$	$\theta-4a$
B6.2	$v_3(B)=3+k,\ v_3(A)=1$	$P_1P_2^2P_3$	$\frac{\theta^3-4a\theta^2}{3}+3$	$\frac{\theta^3-4a\theta^2+6a^2\theta+A}{3}+\theta$	$\theta-4a$

$$TableB11.4: B=s^4+as+b,\ A=4s^3+a,\ as=-4b_3 \pmod{3^{v_3(\triangle)+1}}$$

Case	*Conditions*	$3\mathbb{Z}_K$	β_4	β_3	β_2	β_1
1	$v_3(b) \geq 2,\ a^2 \neq 7 \pmod 9$ $v_3(a^4-a^2+b)=2$	$P_1P_2^2P_3$		$\frac{\theta^3-4a\theta^2}{3}$	$\frac{\theta^3-4a\theta^2+6sa^2\theta+A}{3}+\theta^2$	$\theta-4a$
2	$v_3(b) \geq 2,\ a^2 \neq 7 \pmod 9$, $v_3(a^4-a^2+b) \geq 3$	$P_1P_2^2P_3$		$\frac{\theta^3-4a\theta^2}{3}+3$	$\frac{\theta^3-4a\theta^2+6a^2\theta+A}{3}+\theta$	$\theta-4a$
3	$v_3(b) \geq 2,\ a^2 = 7 \pmod 9$, $v_3(a^4-a^2+b)=2$	$P_1P_2^3$			$\frac{\theta^3+4s\theta^2}{3}+\theta$	$\theta+4s$
4	$v_3(\triangle)=2r+1,\ r\geq 4$	$P_1P_2P_3^2$		$\frac{\theta^3+4s\theta^2+6s^2\theta+A}{3^{r-1}}+\frac{\theta^2+4s\theta}{3}$	$\frac{\theta^3+4s\theta^2}{3}+\theta$	$\theta+4s$
5	$v_3(\triangle)=7$,	$P_1P_2P_3^2$		$\frac{\theta^3+4s\theta^2+6s^2\theta+A}{3^{r-1}}+\frac{\theta^2+4s\theta}{3}$	$\frac{\theta^3+4s\theta^2}{3}+\theta^2$	$\theta+4s$
6	$v_3(\triangle)=2r,\ r\geq 5$ $(\frac{-2B_3}{3})=-1$	$P_1P_2P_3$		$\frac{\theta^2+4s\theta}{3}+\theta$	$\frac{\theta^2+4s\theta+6s^2}{3}+\theta$	$\theta+4s$
7	$v_3(\triangle)=2r\geq 8$ $(\frac{-2B_3}{3})=1$	$P_1P_2P_3P_4$	$\frac{\theta^3+4s\theta^2+6s^2(\theta+3^{r-2}t)}{3^{r-1}}$ $2s^2t^2+B_3=3 \pmod 9$	$\frac{\theta^3+4s\theta^2+6s^2(\theta-3^{r-2}t)}{3^{r-1}}$	$\frac{\theta^2+4s\theta+6s^2}{3}+\theta$	$\theta+4s$

If ($b=7 \pmod 8$, $a=0 \pmod 8$ *and* $1+b+a=0 \pmod{16}$*) or* $b=3 \pmod 8$ *and* $a=4 \pmod 8$*, then let* $s\in\mathbb{Z}$ *such that* $P(X)$ *is* $(X+s)$*-regular, and let* $A=4s+a$, $B=s^4+as+b$ *and* $\theta=\alpha-s$.

$TableC14: v_2(B)\geq 3, v_2(A)\geq 2$

Case	*Conditions*	$2\mathbb{Z}_K$	β_3	β_2	β_1
1	$v_2(B)\geq 2$ $v_2(A)=1$	$P_1^3P_2$		$\frac{\theta^3}{2}+2$	$\frac{\theta^3+A}{2}+\theta$
2	$v_2(B)=2r-1$ $v_2(A)=r\geq 2$	$P_1^2P_2$		$\frac{\theta^2}{2}+2$	$\frac{\theta^2+6}{2}+\theta$
3	$v_2(B)>2r$ $v_2(A)=r\geq 2$	$P_1^2P_2P_3$	$\frac{\theta^3+4s\theta^2+6s^2\theta}{2^r}$	$\frac{(\theta^2+4s\theta+6s^2)(\theta+2^{r-1}t)}{2^r}+2$ $A_2t=1 \pmod 4$	$\frac{\theta^2+6}{2}+\theta$
4	$v_2(B)=2r$ $v_2(A)=r\geq 2$	$P_1^2P_2P_3$	$\frac{\theta^3+4s\theta^2+6s^2\theta}{2^r}$	$\frac{(\theta^2+4s\theta+6s^2)(\theta+2^{r-1}t)}{2^r}$ $A_2t=3 \pmod 4$	$\frac{\theta^2+6}{2}+\theta$
5	$v_2(B)=2r$ $v_2(A)\geq r+1$	$P_1^2P_2^2$		$\frac{\theta^2}{2}+\frac{\theta^3+4s\theta^2+6s^2\theta+A}{2^r}$	$\frac{\theta^2+6}{2}+\theta$

$TableD8: p\geq 5,\ v_p(ab)=0 and v_p(\triangle)\geq 2$

Let $s \in \mathbb{Z}$ *such that* $3as + 4b = 0 \pmod{p^{v_p(\triangle)+1}}$, $\theta = \alpha - s$, $B = s^4 + as + b$ *and* $A = 4s^3 + a$.

Conditions	$p\mathbb{Z}_K$	β_4	β_3	β_2	β_1
$v_p(\triangle) = 2r+1$ $(\frac{-2}{p}) = 1$	$P_1P_2P_3^2$		$\frac{\theta^3+4s\theta^2+6s^2\theta}{p^r}$ $B_p = 6s^2u \pmod p$	$\theta - v_1$ $u^2+2 = p \pmod{p^2}$ $v_1 = -s(2+u) \pmod{p^2}$	$\theta - v_2$ $u^2+2 = p \pmod{p^2}$ $v_2 = -s(2-u) \pmod{p^2}$
$v_p(\triangle) = 2r+1$ $(\frac{-2}{p}) = -1$	$P_1P_2^2$			$\frac{\theta^3+4s\theta^2+6s^2\theta}{p^r}$ $B_p = 6s^2u \pmod p$	$\theta^2 + 4s\theta + 6s^2 + p$
$v_p(\triangle) = 2r$ $(\frac{-2}{p}) = 1$ $(\frac{-6B_p}{p}) = -1$	$P_1P_2P_3$		$\frac{\theta^3+4s\theta^2+6s^2\theta}{p^{r-1}}$ $v_p(6s^2 + B_p) = 1$	$\theta - v_1$ $u^2+2 = p \pmod{p^2}$ $v_1 = -s(2+u) \pmod{p^2}$	$\theta - v_2$ $u^2+2 = p \pmod{p^2}$ $v_2 = -s(2-u) \pmod{p^2}$
$v_p(\triangle) = 2$ $(\frac{-2}{p}) = 1$ $(\frac{-6B_p}{p}) = 1$	$P_1P_2P_3P_4$	$\frac{(\theta^2+4s\theta+6s^2)(\theta+pt)}{p} + \theta^2$ $v_p(6s^2t^2 + B_p) \geq 2$	$\frac{(\theta^2+4s\theta+6s^2)(\theta-pt)}{p} + \theta^2$	$\theta - v_1$ $u^2+2 = p \pmod{p^2}$ $v_1 = -s(2+u) \pmod{p^2}$	$\theta - v_2$ $u^2+2 = p \pmod{p^2}$ $v_2 = -s(2-u) \pmod{p^2}$
$v_p(\triangle) = 2r \geq 4$ $(\frac{-2}{p}) = 1$ $(\frac{-6B_p}{p}) = 1$	$P_1P_2P_3P_4$	$\frac{(\theta^2+4s\theta+6s^2)(\theta+p^rt)}{p^r} + \theta$ $v_p(6s^2t^2 + B_p) = 1$	$\frac{(\theta^2+4s\theta+6s^2)(\theta-p^rt)}{p^r} + \theta$	$\theta - v_1$ $u^2+2 = p \pmod{p^2}$ $v_1 = -s(2+u) \pmod{p^2}$	$\theta - v_2$ $v_p(u^2+2) = 1$ $v_p(v_2 + s(2-u)) \geq 2$
$v_p(\triangle) = 2r$ $(\frac{-2}{p}) = -1$ $(\frac{-6B_p}{p}) = -1$	P_1P_2			$\frac{\theta^3+4s\theta^2+6s^2\theta}{p^{r-1}}$	$\theta^2 + 4s\theta + 6s^2 + p$
$v_p(\triangle) = 2$ $(\frac{-2}{p}) = -1$ $(\frac{-6B_p}{p}) = 1$	$P_1P_2P_3$		$\frac{(\theta^2+4s\theta+6s^2)(\theta+pt)}{p} + \theta^2$ $v_p(6s^2t^2 + B_p) \geq 2$	$\frac{(\theta^2+4s\theta+6s^2)(\theta-pt)}{p} + \theta^2$	$\theta^2 + 4s\theta + 6s^2 + p$ $v_p(6s^2u + B_p) = 1$
$v_p(\triangle) = 2r \geq 4$ $(\frac{-2}{p}) = -1$ $(\frac{-6B_p}{p}) = 1$	$P_1P_2P_3$		$\frac{(\theta^2+4s\theta+6s^2)(\theta+p^rt)}{p^r} + \theta$ $v_p(6s^2t^2 + B_p) = 1$	$\frac{(\theta^2+4s\theta+6s^2)(\theta-p^rt)}{p^r} + \theta$	$\theta^2 + 4s\theta + 6s^2 + p$

Proof of Theorem. All cases, except ($p = 2$, $v_p(b) = 2$ and $v_p(a) \geq 2$), correspond to a situation where $P(X)$ is p-regular. The case : $p = 2$, $v_p(b) = 2$ and $v_p(a) \geq 2$ is handled in $TableA8$ by using technics of Newton polygons of second order.
Denote $C(X)$ the minimal polynomial of a possible ϕ such that $v_p([\mathbb{Z}_K : \mathbb{Z}[\phi]]) = 0$.
$v_p(a) \geq 1$ **and** $v_p(b) \geq 1$**.**

1. $v_p(a) \geq 3$ and $v_p(b) = 3$. Then $\bar{P}(X) = X^4 \pmod p$ and the X-Newton polygon of $P(X)$ is one side of slope $3/4$. Thus, $p\mathbb{Z}_K = P^4$. Since $v_P(\alpha) = 4 \times 3/4 = 3$, then $v_P(\frac{\alpha^3}{p^2}) = 1$ and $P = (p, \frac{\alpha^3}{p^2})$.
2. Let $\phi = \frac{\alpha^3}{p^h}$. Then $C(X) = X^4 + 3\frac{a}{p^h}X^3 + 3\frac{a^2}{p^{2h}}X^2 + \frac{a^3}{p^{3h}}X + \frac{b^3}{p^{4h}}$. It follows that if $v_p(a) = 1$ and $v_p(b) \geq 2$, then $\phi = \frac{\alpha^3}{p}$ is a p-generator of $\mathbb{Z}_K$, $\bar{C}(X) = X(X + a_p)^3 \pmod p$ and $p\mathbb{Z}_K = (p, \alpha)(p, \alpha + a_p)^3$. If $v_p(a) = 2$ and $v_p(b) \geq 3$, then $\phi = \frac{\alpha^3}{p^2}$ is a p-generator of $\mathbb{Z}_K$, $\bar{C}(X) = X(X + a_p)^3 \pmod p$ and $p\mathbb{Z}_K = (p, \phi)(p, \phi + a_p)^3$.
3. If $v_p(b) = 1$ and $v_p(a) \geq 1$, then $v_p(ind(P)) = 0$, then α is a p-generator of $\mathbb{Z}_K$ and $p\mathbb{Z}_K = (p, \alpha)^4$.
4. If $v_p(b) = 2$ and $v_p(a) \geq 2$, then $N_X(P) = S$ is one side such that $P_S(y) = Y^2 + b_p$. It follows that:
   (a) If $(\frac{-b_p}{p}) = -1$, then $p\mathbb{Z}_K = P^2$ such that $v_P(\alpha) = 1$. Thus, $P = (p, \frac{\alpha^3}{p})$.
   (b) If $(\frac{-b_p}{p}) = 1$ and $v_p(a) = 2$, then $p\mathbb{Z}_K = P_1^2P_2^2$. For $\phi = \frac{\alpha^2}{p}$, we have $C(X) = X^4 - Xpa_p^2 + 2b_pX^2 + b_p^2$ and $\bar{C}(X) = (X^2 + b_p)^2 \pmod p$. Let $t \in \mathbb{Z}$ such that $v_p(t^2 + b_p) = 1$. Let $\phi(X) = X + t$ and $\phi(X)^4 - 4t\phi(X)^3 + (6t^2 + 2B)\phi(X)^2 + (-pa_p^2 - 4tb_p - 4t^3)\phi(X) + (b_p^2 + tpa_p^2 + 2t^2b_p + t^4)$ be the $\phi(X)$-adic development

of $C(X)$. Since $v_p(b_p^2+tpa_p^2+2t^2b_p+t^4)=1$, $\frac{\alpha^2}{p}$ is a p-generator of $\mathbb{Z}_K$. As $\bar{C}(X)=(X^2+b_p)^2 \pmod p$, we have $P_1=(p,t+\frac{\alpha^2}{p})$ and $P_2=(p,-t+\frac{\alpha^2}{p})$ $v_p(t^2+b_p)=1$.

(c) If $(\frac{-b_p}{p})=1$ and $v_p(a)=\geq 3$, then $p\mathbb{Z}_K=P_1^2P_2^2$. Let $\phi=\frac{\alpha^3+\alpha^2}{p}$. Since $\alpha^3+\alpha^2\notin\mathbb{Q}$, $K=\mathbb{Q}[\alpha^3+\alpha^2]$ and $\bar{C}(X)=X^4+2b_pX^2-4pb_p^2X+b_p^2 \pmod{p^2}$ and $\bar{C}(X)=(X^2+b_p)^2 \pmod p$. Let $\phi(X)=X+t$ such that $t^2+b_p=0 \pmod p$ and $C(X)=\phi^4(X)-4t\phi^3(X)+(6t^2+2b_p)\phi^2(X)+(-4pb_p^2-4tb_p-4t^3)\phi(X)+(b_p^2+4tpb_p^2+2t^2b_p+t^4)$ be the $\phi(X)$-adic development of $C(X)$. Since $v_p(b_p^2+4tpb_p^2+2t^2b_p+t^4)=1$, $\frac{\alpha^3+\alpha^2}{p}$ is a p-generator of $\mathbb{Z}_K$, $P_1=(p,t+\frac{\alpha^3+\alpha^2}{p})$ and $P_2=(p,-t+\frac{\alpha^3+\alpha^2}{p})$.

(5) $v_2(b)=2$ and $v_2(a)\geq 2$. In that case, using technics of Newton polygons of second order, for every subcase, a monic polynomial $\phi_2\in\mathbb{Z}[X]$ such that $t=(X,1/2,\phi_2)$ is $P(X)$-complet is given (see [5, Def 3.9, p 38]). Let N_2 be the ϕ_2-Newton polygon of second order of $P(X)$. From [5, Cor 3.8, p 38], if N_2 is one side, then $2\mathbb{Z}_K=P^e$, where $e=2e_2$ and e_2 is the ramification index of N_2. If N_2 is two sides, then for every side, $e_2=1$ and then $2\mathbb{Z}_K=P_1^2P_2^2$.

(a) If $v_2(a)=2$, then for $\phi_2=X^2+2$ and N_2 is one side. Thus, $2\mathbb{Z}_K=P^4$, $v_P(\alpha^2+2)=5$ and $2\mathbb{Z}_K=(2,\frac{\alpha^2+2}{2})$.

(b) If $v_2(a)=3$ and $b=4 \pmod{16}$ (resp. $v_2(a)=3$ and $b=12 \pmod{16}$), then for $\phi_2=X^2+2X+2$ (resp. $\phi_2=X^2+2$) and N_2 is one side. Then $2\mathbb{Z}_K=P^4$, $v_P(\alpha^2+2\alpha+2)=7$ (resp. $v_P(\alpha^2+2)=7$) and $2\mathbb{Z}_K=(2,\frac{\alpha^3+2\alpha^2+2\alpha}{4})$ (resp. $2\mathbb{Z}_K=(2,\frac{\alpha^3+2\alpha}{4})$).

(c) If $v_2(a)=4$ and $b=4 \pmod{32}$, then for $\phi_2=X^2+2X+2$, N_2 is one side, $2\mathbb{Z}_K=P^4$ and $v_P(\alpha^2+2\alpha+2)=9$. Thus, $2\mathbb{Z}_K=(2,\frac{\alpha^2+2\alpha+2}{4})$.

(d) If $v_2(a)=4$ and $b=20 \pmod{64}$, then for $\phi_2=X^2+2X-2$, N_2 is one side, $2\mathbb{Z}_K=P^2$ and $v_P(\alpha^2+2\alpha-2)=5$. Thus, $2\mathbb{Z}_K=(2,\frac{\alpha^2+2\alpha-2\alpha}{4})$.

(e) If $v_2(a)\geq 4$ and $b=12 \pmod{32}$, then for $\phi_2=X^2+2$, N_2 is one side, $2\mathbb{Z}_K=P^2$ and $v_P(\alpha^2+2)=4$. Thus, $2\mathbb{Z}_K=(2,\frac{\alpha^3+2\alpha}{4})$.

(f) If $v_2(a)\geq 5$ and $b=20 \pmod{32}$, then for $\phi_2=X^2+2X-2$, N_2 is one side, $2\mathbb{Z}_K=P^4$ and $v_P(\alpha^2+2\alpha-2)=9$. Thus, $2\mathbb{Z}_K=(2,\frac{\alpha^2+2\alpha-2\alpha}{4})$.

(g) If $v_2(a)\geq 5$ and $b=36 \pmod{64}$, then for $\phi_2=X^2+2X+2$, N_2 is one side, $2\mathbb{Z}_K=P^2$ and $v_P(\alpha^2+2\alpha+2)=5$. Thus, $2\mathbb{Z}_K=(2,\frac{\alpha^2+2\alpha-2\alpha}{4})$.

(6) $b=4+64B$ and $a=32+64A$. For $\phi=\frac{\alpha^3-2\alpha-4}{8}$, we have $\bar{C}(X)=X^4+2X^3+3X^2+2X+2 \pmod 4$. Thus, $v_2(C(1))=v_2(C(0))=1$, ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K=(2,\phi)^2(2,\phi+1)^2$.

(7) $b=4+64B$ and $a=64A$. For $\phi=\frac{\alpha^3+4\alpha^2-2\alpha+4}{8}$, we have $\bar{C}(X)=X^4+2X^3+3X^2+2X+2 \pmod 4$. It follows that $v_2(C(1))=v_2(C(0))=1$, ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K=(2,\phi)^2(2,\phi+1)^2$.

(8) $b=28+32B$ and $a=16+32A$. For $\phi=\frac{\alpha^2+2}{4}$, we have $C(X)=X^4-2X^3+(5+4B)X^2+(-16A^2-8-16A-4B)X+6+8A+8A^2+8B+4B^2=(X+1)^2X^2 \pmod 2$. Since $v_2(C(0))=v_2(C(1))=1$, we have ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K=(2,\phi+1)^2(2,\phi)^2$.

(9) $b=52+64B$ and $a=16+64A$. For $\phi=\frac{\alpha^3-2\alpha^2-2\alpha}{8}$, $C(X)=X^4+2X^3+3X^2+2X+2 \pmod 4$. Thus, $v_2(C(0))=v_2(C(1))=1$, ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K=(2,\phi+1)^2(2,\phi)^2$.

(10) $b = 52 + 64B$ and $a = 48 + 64A$. For $\phi = \frac{\alpha^3-2\alpha^2+6\alpha}{8}$, $C(X) = X^4 + 2X^3 + 3X^2 + 2X + 2 \pmod 4$. Thus, $v_2(C(0)) = v_2(C(1)) = 1$, ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K = (2, \phi + 1)^2(2, \phi)^2$.

(11) $b = 28 + 32B$ and $a = 32A$. For $\phi = \frac{\alpha^2+12\alpha+2}{4}$, $C(X) = X^4 + 2X^3 + X^2 + 2 \pmod 4$. Thus, $v_2(C(0)) = v_2(C(1)) = 1$, ϕ is a 2-generator of $\mathbb{Z}_K$ and $2\mathbb{Z}_K = (2, \phi+1)^2(2, \phi)^2$.

$v_p(a) = 0$ **and** $v_p(b) \geq 1$**.**
If $p \neq 3$, then $v_p(ind(P)) = 0$, α is a p-genrator of $\mathbb{Z}_K$ and $p\mathbb{Z}_K$ is p-analogous to $\bar{P}(X)$.
For $p = 3$, let $F(X) = P(X-a) = X^4-4aX^3+6a^2X^2+AX+B$ and $\theta = \alpha+a$ ($A = -a(4a^2-1)$ and $B = (a^4 - a^2 + b)$).

(1) If $v_3(B) = 1$, then $v_3(ind(P)) = 0$ and $3\mathbb{Z}_K = (3, \alpha + a)^3(3, \alpha)$.
(2) If $v_3(B) = 2$ and $v_3(A) \geq 2$, then $3\mathbb{Z}_K = P_1P_2^3$, $v_{P_1}(\theta - 4a) = 1$, $v_{P_1}(\theta) = 0$, $v_{P_2}(\theta - 4a) = 0$ and $v_{P_2}(\theta) = 2$. Thus, $P_1 = (3, \theta - 4a)$, $P_2 = (3, \frac{\theta^3-4a\theta^2}{3})$.
(3) If $v_3(A) = 1$ and $v_3(B) \geq 2$, then $3\mathbb{Z}_K = P_1P_2^2P_3$.
Since $\theta^4 - 4a\theta^3 = -6a^2\theta^2 + A\theta + B$ and $v_{P_1}(\theta) = 0$, we have $v_{P_1}(\theta^3 - 4a\theta^2) = 1$ and $P_1 = (3, \theta - 4a)$.
If $v_3(B) = 2$, then let $\beta_3 = \frac{\theta^3-4a\theta^2}{3}$ and $\beta_2 = \frac{\theta^3-4a\theta^2+6sa^2\theta+A}{3} + \theta^2$. Since $v_{P_3}(\theta) = v_{P_2}(\theta) = 1$ and $v_{P_1}(\theta) = 0$, we have $v_{P_3}(\beta_3) = 1$ and $v_{P_i}(\beta_3) = 0$ for $i \neq 3$. We have also, $v_{P_2}(\beta_2) = 1$ and $v_{P_i}(\beta_3) = 0$ for $i \neq 2$.
The case $v_3(B) \geq 3$ is similar to the previous case.
(4) If $v_3(A) \geq 2$ and $v_3(B) \geq 3$ ($v_3(\triangle) \geq 6$), then let $s \in \mathbb{Z}$ such that $as = -4b_3 \pmod{3^{v_3(\triangle)+1}}$.
Let $F(X) = P(X+s) = X^4+4sX^3+6s^2X^2+AX+B$ and $\theta = \alpha-s$. Since $A = 4s^3+a$ and $B = s^4 + as + b$, $v_3(A) = v_3(B) = v_3(\triangle) - 3 \geq 3$. It follows that:
If $v_3(\triangle) = 6$, then for $\phi = \frac{\theta^3+4s\theta^2+6s^2\theta+A}{9}$, we have $\bar{C}(X) = X(X^3 + 2s^2B_3X - 4sB_3^2) \pmod 3$. So, if $B_3 = 1 \pmod 3$, then $\bar{C}(X) = X(X^3 - X - s) \pmod 3$ and $3\mathbb{Z}_K = (3, \phi)(3, \phi^3 - \phi - s)$. If $B_3 = -1 \pmod 3$ and $s = 1 \pmod 3$, then $\bar{C}(X) = X(X^2 - X - 1)(X + 1) \pmod 3$. If $B_3 = -1 \pmod 3$ and $s = -1 \pmod 3$, then $\bar{C}(X) = X(X^2 + X - 1)(X - 1) \pmod 3$
(5) If $v_3(\triangle) \geq 7$, then $N_X(F) = S_3 + S_2 + S_1$, $F_{S_1}(Y) = Y + 4s$ and $F_{S_2}(Y) = sY - 1$.
Since $\theta^3(\theta^2 + 4s) = -(6s^2\theta^2 + A\theta + B)$, then $P_1 = (3, \theta + 4s)$.
  (a) If $v_3(\triangle) = 2r + 1$ ($r \geq 3$), then $v_3(A) = v_3(B) = 2r - 2$, $F_{S_3}(Y) = 2s^2Y + B_3$, $3\mathbb{Z}_K = P_1P_2P_3^2$, $v_{P_3}(\theta) = 2r - 3$, $v_{P_2}(\theta) = 1$ and $v_{P_1}(\theta) = 0$. For every $i \neq j$, $v_{P_i}(\beta_i) = 1$ and $v_{P_i}(\beta_j) = 0$.
  (b) If $v_3(\triangle) = 2r$, then $F_{S_3}(Y) = 6s^2Y^2 + B_3$. Thus, if $(\frac{-2B_3}{3}) = -1$, then $3\mathbb{Z}_K = P_1P_2P_3$. If $(\frac{-2B_3}{3}) = -1$, then $3\mathbb{Z}_K = P_1P_2P_3P_4$.
  If $(\frac{-2B_3}{3}) = -1$, then let $\beta_3 = \frac{\theta^2+4s\theta}{3} + \theta + 3$ and $\beta_2 = \frac{\theta^2+4s\theta+6s^2}{3} + \theta$.
  If $(\frac{-2B_3}{3}) = 1$, then $F_{S_3}(Y) = 6s^2(Y + t)(Y - t)$, where $6s^2t^2 + B_3 = 0 \pmod 3$. Thus, $3\mathbb{Z}_K = P_1P_2P_3P_4$, where P_3 and P_4 are respectively attached to $Y + t$ and $Y - t$. Let $\phi = X + 3^{r-2}t$ and consider the ϕ-adic development of $F(X)$, where $2s^2t^2 + B_3 = 3 \pmod 9$. We have $v_{P_3}(\phi(\theta)) = r - 1$ and $v_{P_4}(\phi(\theta)) = r - 2$. Therefore, $\beta_4 = \frac{\theta^3+4s\theta^2+6s^2(\theta+3^{r-2}t)}{3^{r-1}}$, $\beta_3 = \frac{\theta^3+4s\theta^2+6s^2(\theta-3^{r-2}t)}{3^{r-1}}$ and $\beta_2 = \frac{\theta^2+4s\theta+6s^2}{3} + \theta$, where $2s^2t^2 + B_3 = 3 \pmod 9$ satisfy: $v_{P_3}(\theta^3 + 4s\theta^2) = v_{P_4}(\theta^3 + 4s\theta^2) = 2(r-1)$, $v_{P_3}(\theta + 3^{r-2}t) = r - 1$, $v_{P_4}(\theta + 3^{r-2}t) = r - 2$, $v_{P_3}(\theta - 3^{r-2}t) = r - 2$ and $v_{P_4}(\theta - 3^{r-2}t) = r - 1$. So, $v_{P_i}(\beta_i) = 1$ and $v_{P_j}(\beta_i) = 0$ for every $i \neq j$.

$v_p(a) \geq 1$ **and** $v_p(b) = 0$**.**
If $p \neq 2$, then $v_p(ind(P)) = 0$. Thus, α ia p-generator of $\mathbb{Z}_K$, and then $p\mathbb{Z}_K$ is p-analogous to $\bar{P}(X) = X^4 + aX + b \pmod p$. For $p \geq 5$, if $(\frac{-b}{p})_4 = 1$, then $\bar{P}(X) = (X - u)(X + u)(X^2 + u^2)$, where $u^4 = -b \pmod p$.
$(\frac{-b}{p})_4 \neq 1$, then $\bar{P}(X)$ is irreducible in $\mathbb{F}_p[X]$ or $\bar{P}(X) = (X^2 + rX + s)(X^2 - rX +$

$t)$ (mod p). In the last case, ($r=0$ (mod p), $s=-t$ (mod p) and $s^2=-b$ (mod p)) or ($r\neq 0$ (mod p), $s=t$ (mod p), $2s=r^2$ (mod p) and $s^2=b$ (mod p)); i.e., $\bar{P}(X)=(X^2+s)(X^2-s)$ (mod p), where $s^2=b$ (mod p) or $\bar{P}(X)=(X^2+uX+s)(X^2-uX+s)$ (mod p), where $u^4=4b$ (mod p) and $2s=u^2$ (mod p).
For $p=2$, let $F(X)=P(X+1)=X^4+4x^3+6X^2+AX+B$ and $\theta=\alpha-1$.

(a) If $v_2(B)=3$, $A=4$ (mod 8), then for $\phi=\frac{\theta^3+4\theta^2+6\theta}{4}$, we have $C(X)=X^4+(3+2K)X^3+(2+2L)X^2+(3+2K)X+1+2L$ (mod 4), where $A=4+8K$ and $B=8+16L$. Since $C(1)=2$ (mod 4), ϕ is a 2-generator of $\mathbb{Z}_K$, $2\mathbb{Z}_K=P_1^2P_2$, where $P_1=(2,\phi+1)$ and $P_2=(2,\phi^2+\phi+1)$.

(b) $b=3$ (mod 8) and $a=4$ (mod 8) or ($b=7$ (mod 8), $a=0$ (mod 8) and $1+b+a=0$ (mod 16)), then let $s\in\mathbb{Z}$ such that $P(X)$ is $X+s$-regular. Let $F(X)=P(X-s)=X^4+4sX^3+6s^2X^2+AX+B$ and $\theta=\alpha-s$. It follows that:
    (i) $v_2(A)=1$ and $v_2(B)\geq 2$. Then $2\mathbb{Z}_K=P_1^3P_2$.
    (ii) If $v_2(A)=r$ and $v_2(B)=2r-1$, then $N_X(F)=S_1+S_2$ such that $F_{S_1}(Y)=Y+1$ and $F_{S_2}(Y)=Y^2+Y+1$. Thus, $2\mathbb{Z}_K=P_1^2P_2$, where $v_{P_1}(\theta)=1$ and $v_{P_2}(\theta)=r-1$. Hence, for every $i\neq j$, $v_{P_i}(\beta_i)=1$ and $v_{P_i}(\beta_j)=0$.
    (iii) If $v_2(A)=r$ and $v_2(B)\geq 2r$, then $N_X(F)=S_1+S_2+S_3$ such that $F_{S_1}(Y)=Y+1$ and $F_{S_3}(Y)=F_{S_2}(Y)=Y+1$. Thus, $2\mathbb{Z}_K=P_1^2P_2P_3$. Let $\beta_3=\frac{\theta^3+4s\theta^2+6s^2\theta}{2^r}$, $\beta_2=\frac{(\theta^2+4s\theta+6s^2)(\theta+2^{r-1}t)}{2^r}$ and $\beta_1=\frac{\theta^2+6}{2}+\theta$, where $t\in\mathbb{Z}$ is choused such that $v_{P_2}(\theta+2^{r-1}t)=r$: Let $\phi(X)=X+2^{r-1}t$ and consider the $\phi(X)$-adic of $F(X)$. If $v_2(B)=2r$, then $A_2t=3$ (mod 4) and f $v_2(B)>2r$, then $A_2t=1$ (mod 4).
    (iv) If $v_2(A)=r+1+k$ and $v_2(B)=2r$, then $N_X(F)=S_1+S_2$ such that $F_{S_1}(Y)=Y+1$ and $F_{S_3}(Y)=Y+1$. Thus, $2\mathbb{Z}_K=P_1^2P_2^2$, where $v_{P_1}(\theta)=1$ and $v_{P_2}(\theta)=2r-1$.

$v_p(ab)=0$.
If $p\in\{2,3\}$, then $v_p(ind(P))=0$ and then $p\mathbb{Z}_K$ is p-analogous to $\bar{P}(X)$. For $p\geq 5$, let $s\in\mathbb{Z}$ such that $3as+4b=0$ (mod $p^{v_p(\triangle)+1}$), $\theta=\alpha-s$ and $F(X)=P(X+s)=X^4+4sX^3+6s^2X^2+AX+B$. Then $v_p(A)=v_p(B)=v_p(\triangle)$.
If $v_p(\triangle)=0$, then $p\mathbb{Z}_K$ is p-analogous to $\bar{P}(X)$. If $v_p(\triangle)=1$ and $(\frac{-2}{p})=1$, then $p\mathbb{Z}_K=P_1^2P_2P_3$, where $P_1=(p,\theta)$, $P_2=(p,\theta+u)$, $P_3=(p,\theta-u)$, $\theta=\alpha-s$ and $3at+4b=0$ (mod p^2).
If $v_p(\triangle)=1$ and $(\frac{-2}{p})=-1$, then $p\mathbb{Z}_K=P_1^2P_2$, where $P_1=(p,\theta)$ and $P_2=(p,\theta^2+4s\theta+6s^2)$. If $v_p(\triangle)\geq 2$, then let $s\in\mathbb{Z}$ such that $3as+4b=0$ (mod $p^{v_p(\triangle)+1}$), $\theta=\alpha-s$ and $F(X)=P(X+s)=X^4+4sX^3+6s^2X^2+AX+B$, where $B=s^4+as+b$ and $A=4s^3+a$. Then $v_p(A)=v_p(B)=v_p(\triangle)$ and $N_X(F)=S_0+S_1$ with respective slopes 0 and $\frac{v_p(\triangle)}{2}$. It follows that:

(a) If $v_p(\triangle)=2r+1$, then $N_X(F)=S_1+S_2$ such that $F_{S_1}(Y)=Y^2+4sY+6s^2$ and $F_{S_2}(Y)=6s^2Y+B_p$. Thus, if $(\frac{-2}{p})=1$, then $p\mathbb{Z}_K=P_1P_2P_3^2$. If $(\frac{-2}{p})=-1$, then $p\mathbb{Z}_K=P_1P_2^2$.

(b) If $v_p(\triangle)=2r$, then $N_X(F)=S_1+S_2$ such that $F_{S_1}(Y)=Y^2+4sY+6s^2$ and $F_{S_2}(Y)=6s^2Y^2+B_p$. Thus, $(\frac{-2}{p})=1$ and $(\frac{-6B_p}{p})=-1$, then $p\mathbb{Z}_K=P_1P_2P_3$.
If $(\frac{-2}{p})=-1$ and $(\frac{-6B_p}{p})=-1$, then $p\mathbb{Z}_K=P_1P_2$.
If $(\frac{-2}{p})=-1$ and $(\frac{-6B_p}{p})=1$, then $F_{S_2}(Y)=6s^2(Y-t)(Y+t)$, where t is a root of $F_{S_2}(Y)$ in $\mathbb{F}_3$. Thus, $p\mathbb{Z}_K=P_1P_2P_3$, where P_1, P_2 and P_3 are respectively attached to S_1, $Y+t$ and $Y-t$. Let $\beta_2=\frac{(\theta^2+4s\theta+6s^2)(\theta+p^rt)}{p^r}+\theta$ and

$\beta_3 = \frac{(\theta^2+4s\theta+6s^2)(\theta-p^rt)}{p^r}+\theta$, where $t \in \mathbb{Z}$ is choused such that $v_{P_2}(\theta+p^rt) = r+1$, $v_{P_3}(\theta+p^rt) = r$, $v_{P_2}(\theta+-p^rt) = r$ and $v_{P_3}(\theta-p^rt) = r+1$: If $r \geq 2$, then $6s^2t^2+B_p = p \pmod{p^2}$. If $r = 1$, then $6s^2t^2+B_p = 0 \pmod{p^2}$. In that way, we have $P_2 = (p, \frac{(\theta^2+4s\theta+6s^2)(\theta+p^rt)}{p^r}+\theta)$ and $P_3 = (p, \frac{(\theta^2+4s\theta+6s^2)(\theta-p^rt)}{p^r}+\theta)$. If $(\frac{-2}{p}) = 1$ and $(\frac{-6B_p}{p}) = 1$, then $p\mathbb{Z}_K = P_1P_2P_3P_4$.

Examples 2.5. *Let* $P(X) = X^4+aX+b \in \mathbb{Z}[X]$ *be an irreducible polynomial,* α *a complex root of* $P(X)$ *and* $K = \mathbb{Q}[\alpha]$.

(1) $a = 2^{10}.5$ *and* $b = 2^9.3.5$. *Let* $\theta = \frac{\alpha}{4}$. *Then* θ *is integral with minimal polynomial* $F(X) = X^4+80X+30$ *and* $K = \mathbb{Q}[\theta]$. *Thus* $2\mathbb{Z}_K = (2,\theta)^4$, $3\mathbb{Z}_K = (3,\theta)(3,\theta-1)^3$, $5\mathbb{Z}_K = (5,\theta)^4$ *and for every* $p \notin \{2,3,5\}$, $p\mathbb{Z}_K$ *is* p*-analogous to* $\bar{F}(X)$.

(2) $a = 48$ *and* $b = 188$. *From* $TableA8$, *row* $A8.9$, *we have* $2\mathbb{Z}_K = (2, \frac{\alpha^2+2}{4})^2(2, \frac{\alpha^2+6}{4})^2$.

(3) $a = 144$ *and* $b = 36$. *From* $TableA8$, *row* $A8.4$, *we have* $2\mathbb{Z}_K = (2, \frac{\alpha^2+2\alpha+2}{4})^4$. *Since* $(\frac{-b_3}{3}) = -1$, *from* $TableA$, *row* $A4$ $3\mathbb{Z}_K = P^2$, *where* $P = (3, \frac{\alpha^3}{3})$. *Since* $\triangle = -2^{14}.3^6.971$, *for every* $p \notin \{2,3,5\}$, $p\mathbb{Z}_K$ *is* p*-analogous to* $\bar{P}(X)$.

(4) $a = 28$, $b = 189$ *and* $p = 2$. *Let* $F(X) = P(X+1) = X^4+4X^3+6X^2+32X+218$. *Since* $B = 218 = 2.109$, *we have* $2\mathbb{Z}_K = (2,\alpha-1)^4$.

(5) $a = 22$ *and* $b = 66$. *Then* $\triangle = 2^4.3^5.11^3.13$. *We have* $2\mathbb{Z}_K = (2,\alpha)^4$, $11\mathbb{Z}_K = (11,\alpha)^4$, $13\mathbb{Z}_K = (13,\alpha)(13,\alpha^3+9)$ *and for every prime* $p \notin \{2,3,11,13\}$, $p\mathbb{Z}_K$ *is* p*-analogous to* $\bar{P}(X)$. *For* $p = 3$, *since* $a^2 = 7 \pmod 9$, $b = 3 \pmod 9$ *and* $b+a^4-a^2 = 9.25982$, *from* $TableB$, *row* $B10$, $3\mathbb{Z}_K = P_1P_2^3$, *where* $P_1 = (3,\alpha-5a)$ *and* $P_2 = (3, \frac{\theta^3-4a\theta^2}{3})$.

(6) $a = 3^6.5^5.139$ *and* $b = 2^2.3^5.5^5.139$. *Let* $\theta = \frac{\alpha}{15}$. *Then* θ *is integral with minimal polynomial* $F(X) = X^4+AX+B$ *and* $K = \mathbb{Q}[\theta]$, *where* $A = 3^3.5^2.139$ *and* $B = 2^2.3.5.139$. *Thus* $2\mathbb{Z}_K = (2,\theta+1)(2,\theta^2+\theta+1)(2,\theta)$, $3\mathbb{Z}_K = (3,\theta)^4$, $5\mathbb{Z}_K = (5,\theta)^4$, $139\mathbb{Z}_K = (139,\theta)^4$. *Since* $\triangle = 2092367789117959822875 = 202317851.7.3^3.5^3.139^3.163$, *for every* $p \notin \{3,5,139\}$, $p\mathbb{Z}_K$ *is* p*-analogous to* $\bar{F}(X)$.

References

[1] S. Alaca and K. S. Williams, p-integral basis of a quartic field defined by a trinomial X^4+aX+b, Far East J. Math. Sci. 12 (2004), 137-168.

[2] Blair K. Spearman and Knneth S. Williams, The prime ideal factorization of 2 in pure quartic fields with index 2, Math. J. Okayama Univ 48(2006), 43-46.

[3] H. Cohen, A course in computational algebraic number theory, GTM 138, Springer-Verlag Berlin Heidelberg, New York, Paris, Tokyo, second correction (1995).

[4] J. Guardia, J. Montes and E. Nart, Newton Polygons of Higher Order in Algebraic Number Theory, arxiv.org/abs/0807.2620v2 [Math.NT] 31 October 2008.

[5] J. Guardia, J. Montes and E. Nart, Higher Newton polygons in the computation of discriminant and prime ideal decomposition in number fields, arXiv:0807.4065v3 [math.NT] 3 November 2008.

[6] P. LLorente, E. Nart and N. Vila, Decompositin of primes in number fields defined by trinomials, J. Theorie des nombres de Bourdeaux, tome 1, 1(1991), 27-41.

Lhoussain El Fadil FPO, P.O. Box 638-Ouarzazte 45000, Morocco
lhouelfadil@hotmail.com